

DETERMINATIONS OF ENDOCRINE DISRUPTORS IN ENVIRONMENTAL USING SPE AND GAS CHROMATOGRAPHY- TANDEM MASS SPECTROMETRY

José Vera, *REQUIMTE, Faculdade de Ciências da Universidade do Porto, Portugal and REQUIMTE, Instituto Superior de Engenharia do Porto, Portugal*

Valentina F. Domingues, *REQUIMTE, Instituto Superior de Engenharia do Porto, Portugal*

Cristina Delerue-Matos, *REQUIMTE, Instituto Superior de Engenharia do Porto, Portugal*

Background and Aims: Endocrine disrupting compounds (EDCs) include a wide variety of pollutants, namely substances of different sources of contamination: domestic, industrial and agricultural e.g. pesticides estrogens, heavy metals and plasticizers. The ECDs are present in the environment at low levels of concentration and they can only be determined by using a preconcentration technique¹. Gas chromatography/mass spectrometry (GC-MS) is commonly accepted as one of the most powerful techniques for the separation, identification and quantification of EDCs, even in the nanogram range².

Methods: Chromatographic analyses were carried out in a Thermo GC ULTRA Gas Chromatograph Mass Spectrometer equipped with a ZB-XLB. High-purity helium at a constant flow rate of 1.3mL/min was used as carrier gas. An AI/AS 3000 auto injector was used. Injections (1µL) were made in the split/splitless mode. Samples were analyzed using the following oven temperature programme: initial temperature 60 C (held for 1 min), increased by 20 C/min to 200 C (held for 1 min), increased again by 5°C/min to 245°C and held at this temperature for 40 min.

Results: We have optimized the method by determining the following ECDs; Atrazine-desethyl, HCB, Atrazine-deuterated, Atrazine, Simazine, Lindane, Vinclozoline, Alaclhor, Malathion, Aldrin, α-Endosulfan, p,p'-DDE, Dieldrin, DDD, Endrin, o,p'-DDT, β-Endosulfan, Methoxychlor, Bifenthrin, Fenproprathrin, Iprodione, λ-Cyhalothrin, Permethrin, β-Cyfluthrin, Cypermethrin, Fenvalerate and Deltamethrin all present good linearity.

Conclusions: The limit of detection achieved allows the determination above the legal limits.

References:

Brossa, L.; Marcé, R. M.; Borrull, F.; Pocurull, E., Determination of endocrine-disrupting compounds in water samples by on-line solid-phase extraction-programmed-temperature vaporisation-gas chromatography-mass spectrometry. *Journal of Chromatography A* 2003, 998 (1-2), 41-50.

Mansilha, C.; Melo, A.; Rebelo, H.; Ferreira, I. M. P. L. V. O.; Pinho, O.; Domingues, V.; Pinho, C.; Gameiro, P., Quantification of endocrine disruptors and pesticides in water by gas chromatography-tandem mass spectrometry. Method validation using weighted linear regression schemes. *Journal of Chromatography A* 2010, 1217 (43), 6681-6691.